



# Search for a new X boson and Dark Photons in NA64 at the CERN SPS.



DSU-2018  
June 2018 Annecy




# Outline

- Motivation
- The NA64 experiment
- Runs NA64
- Simulation of the Dark Matter production
- Analysis of the data
- Results on  $A'$  in invisible mode
- Plans for the invisible mode
- Visible mode: X-boson, motivation
- Event selection, efficiency, backgrounds
- Results on the X-boson search
- Conclusion, near and more distant plans of NA64

# Vector portal to Dark Sector


Okun, Holdom' 86 ..  $\alpha_D = e_D^2/4\pi$



- new massive boson  $A'$  (dark photon) which has kinetic mixing with ordinary photon:  $\Delta L = \epsilon/2 F^{\mu\nu} A'_{\mu\nu}$
- Production:  $A'$  - bremsstrahlung  $e^- Z \rightarrow e^- Z A'$ ,  $\sigma \sim Z^2 \epsilon^2 / m_{A'}^2$
- Decays:
  - Visible:  $A' \rightarrow e^+e^-, \mu^+\mu^-, \text{hadrons}, \dots$
  - Invisible:  $A' \rightarrow \chi\chi$  if  $m_{A'} > 2m_\chi$  assuming  $\alpha_{DM} \sim \alpha \gg \epsilon$ .  
Can explain  $(g-2)_\mu$ , astrophys. observations
- Cross section for  $\chi$ -DM annihilation:  $\sigma v \sim [\alpha_{DM} \epsilon^2 (m_\chi/m_{A'})^4] \alpha / m_\chi^2$



# NA64 experiment setup (invisible mode)



~50 researchers from 12 institutes


Proposed in 2014, first test runs in 2015



# NA64 experiment setup




# Search for $A' \rightarrow$ invisible decays at CERN SPS



S.Andreas et al., arXiv: 1312.3309  
S.G., PRD(2014)

## Main components :

- clean 100 GeV e- beam
- e- tagging system: tracker+SRD
- fully hermetic ECAL+ HCAL



$$SES < 10^{-11}/EOT$$

## Signature:

- in: 100 GeV e- track
- out:  $E_{ECAL} < E_0$  shower in ECAL
- no energy in Veto and HCAL

## Background:

- ◆  $\mu, \pi, K$  decays in flight
- ◆ Tail  $< 50$  GeV in the e- beam
- ◆ Energy leak from ECAL+HCAL



# Summary of the NA64 runs


- **First run 12.10-09.11 2016, 4 w, invisible mode configuration**
  - 23 October → start data taking;
  - Subrun1 EOT  $\sim 2 \times 10^{10}$ ,  $S_0$  rate  $1.5 \div 2.2 \times 10^6$ ;
  - Subrun2 EOT  $\sim 1.5 \times 10^{10}$ ,  $S_0$  rate  $2.4 \div 3.2 \times 10^6$ ;
  - Subrun3 EOT  $\sim 1.0 \times 10^{10}$ ,  $S_0$  rate  $4.6 \div 5.0 \times 10^6$ ;  **$\sim 0.6$  day**
  - **Total number  $\sim 4.5 \times 10^{10}$  eot**
  
- **Second run 09.09-01.10 2017, from 22.09 visible mode configuration**
  - Subrun 1 WCAL 40X0 EOT  $\sim 2.4 \times 10^{10}$ ,  $S_0$  rate  $\sim 3 \times 10^6$ ;
  - Subrun 2 WCAL 30X0 EOT  $\sim 3 \times 10^{10}$ ,  $S_0$  rate  $4-5 \times 10^6$ ;
  - **Total EOT  $\sim 5.4 \times 10^{10}$**



# Simulation of $eZ \rightarrow eZA'$ ; $A' \rightarrow$ invisible @ BG

GEANT4 + code for  $A'$  emission in the process of e-m shower development.  $\sigma(eZ \rightarrow eZA')$  from Bjorken et al. 2009

Gninenko, Kirsanov,  
Krasnikov, Kirpichnikov  
PRD(2016)






# Simulation of $eZ \rightarrow eZA'$

- The signal process is simulated using simplified Weizsaecker – Williams (WW) approximation (Bjorken et al., 2009)
- **More exact calculations that use the full matrix element** were performed recently (2016, 2017) (arXiv:1712.05706 [hep-ph], accepted in Phys. Lett. B)
- We started to use these calculations this year
- They are implemented as **K-factors** to the total cross section. The latter can be decreased by as much as factor 15 w.r.t. the simplified WW approximation at  $M_A \sim 1$  GeV
- The differential cross section (essentially the distribution of the energy fraction transferred to  $A'$ ) from WW is used. The difference is small because both WW and exact are strongly peaked near 1. The  $A'$  spectrum is determined mainly by the EM shower development




# K-factors to $eZ \rightarrow eZA'$






# Reconstruction: key moments


- Synchrotron Radiation detectors (SRD) made as lead – scintillator sandwiches suppress pions and other particles heavier than electrons that are present in the beam by a factor of  $10^{-5}$
- The shower profile in ECAL is compared to the profile of true electrons in order to further suppress wrong particles.
- Micromegas track detectors are used to reconstruct the momentum of electron before the ECAL in order to suppress small fraction of soft electrons from interactions on beam line elements.



# Dimuon production as a reference process



- There is an excellent reference process: **gamma to muons conversion**. It is rather rare and has many similarities with our signal
- Several  $10^4$  dimuon pairs with both muons reaching all HCAL modules are registered in the 2016 runs
- The process is available in GEANT4, off by default
- We bias the cross section in GEANT4 by a factor of 200 in order to have good statistics with reasonable CPU time.
- Good agreement DATA - MC





# Dimuon reconstruction


## HCAL module 3




Dimuons selection:  $E_{\text{ECAL}} < 60 \text{ GeV}$   
 $2.5 < E_{\text{HCAL1}} < 6.35$   
 $2 < E_{\text{HCAL3}} < 6.35$

Left plot: number of dimuons in DATA  $\sim 0.92$  of MC prediction  $\rightarrow$  efficiency correction


# Background






# Background


- As mentioned above, the sources of background are decays in flight and various impurities of the beam (softer electrons etc.)
- The BG from decays was estimated by biasing the life times in GEANT4
- The second BG is higher and difficult to simulate. We estimated it using extrapolation from the “side bin” , i.e. from what we see beside our “signal box” preliminarily defined as “ $E_{\text{ECAL}} < 50 \text{ GeV}$ ”



# Background: example of extrapolation




Total predicted background  $\sim 0.17$




# Analysis: efficiency corrections and uncertainties

Efficiency type	Method	Efficiency	uncertainty
Trigger and SRD selection, DAQ	Dimuons analysis	0.91	10%
VETO cut	Comparison MC - data in calib. runs	1	5%
HCAL cut	Comparison MC - data in calib. runs	0.99	5%

Veto: cut at 0.01 GeV




HCAL0: cut at 1 GeV






# Analysis

- Data collected in the autumn 2016 run are divided in 3 bins: low, medium and high intensity
- For each bin the background, efficiency corrections and their uncertainties are estimated
- The expected sensitivity was calculated with ProfileLikelihood method
- The limits are calculated with  $CL_S$  method





# Analysis: optimization



The optimization confirmed the preliminary choice of the  $E_{\text{ECAL}}$  cut: 50 GeV

# Results



arXiv:1710.00971 [hep-ph]  
Phys. Rev. D 97, 072002 (2018)




# Future plans

- More data in invisible mode are collected in 2017 and 2018
- The analysis of these data is now been performed
- New results are expected this and next year




# Search for a new X-boson decaying to $e^+e^-$



# ATOMKI $^8\text{Be}^*$ anomaly: a new 17 MeV gauge boson?

$^7\text{Li}(p,\gamma)^8\text{Be}$ ,  $M_X = 17 \text{ MeV}$



Feng et al, 2016





FIG. 5. Invariant mass distribution derived for the 18.15 MeV transition in  $^8\text{Be}$ .


X cannot be A' due to constraints from  $\pi^0$ -

$\rightarrow X\gamma$  decay:




$$\Gamma(\pi^0 \rightarrow X\gamma) \sim (\epsilon_u q_u - \epsilon_d q_d)^2 \sim 0$$


if  $2\epsilon_u = -\epsilon_d \rightarrow$  **protophobic X**



# Search for the $X(A') \rightarrow e^+e^-$ decays




- X decays outside WCAL dump
- **Signature:** two separated showers from a single  $e^-$ 
  - $E_{WC} < E_0$ , and  $E_0 = E_{WC} + E_{EC}$
  - $\theta_{e^+e^-}$  too small to be resolved
    - background
      - Beam hadrons
  - SRD  $e^-$ -tagging is a key point






# Event selection: criteria

- SRD tag
- $E_{\text{WCAL}} < 70 \text{ GeV}$  (preliminary trigger selection  
 $E_{\text{WCAL}} < \sim 75 \text{ GeV}$ )
- $E_{\text{V2}} < 0.6 \text{ MIP}$  (no charged particles after WCAL).
- $E_{\text{S4}} > 1.5 \text{ MIP}$  (two charged particles in ECAL).  
Control region for neutrals:  $E_{\text{S4}} < 0.7 \text{ MIP}$
- $E_{\text{WCAL}} + E_{\text{ECAL}} > 85 \text{ GeV}$
- Shower profile in ECAL compatible with electron (or with two very close electrons)
- Small energy in VETO and HCAL




# Checks of efficiency to signal

- **Dimuons (gamma to muons conversion)** are used also in this configuration: efficiency corrections
- **Electron calibration runs** are used to compare the the distributions in the detectors used as veto: V2, VETO, HCAL
- Checking the shower profile. We cannot have a single electron in ECAL in this configuration. We selected **muons** from the hadron calibration runs that emit **hard delta electron in ECAL**. We require  $EECAL > 20$  GeV and we select events with small activity in ECAL, VETO. All such events have  $\chi^2$  below our cut.



Main background from  $K^0_S \rightarrow \pi^0 \pi^0 \rightarrow \gamma' s \rightarrow e^+e^-$  decay chain



We used two **control samples** to estimate this BG: **fully neutral events** with and without cut on  $E_{\text{HCAL}}$ .

Two methods to estimate this background:


First: sample with removed cut on  $E_{\text{HCAL}}$ . Main contribution from neutrons, also  $K^0_L$  contribute

Second: sample with cut on  $E_{\text{HCAL}}$

**Method I:** selection of neutral hadronic final state:  $n:K^0 \sim 10:1 \Rightarrow n_{K^0} \sim 10^2 K^0$  **Method II:** selection of e.m. neutrals ( $\gamma'$  s from  $K^0_S$  chain)  $\Rightarrow n_{K^0} \sim 1.5 \times 10^2 K^0$

**Consistent estimates of  $K^0_S$**

Then use Geant4 MC to estimate the number of  $K^0_S$  events with conversion before S4




## Final estimate of the background

Source of background	Events
$e^+e^-$ pair production by punchthrough $\gamma$	$< 0.001$
$K_S^0 \rightarrow 2\pi^0; \pi^0 \rightarrow \gamma e^+e^-$ or $\gamma \rightarrow e^+e^-; K_S^0 \rightarrow \pi^+\pi^-$	$0.06 \pm 0.034$
$\pi N \rightarrow (\geq 1)\pi^0 + n + \dots; \pi^0 \rightarrow \gamma e^+e^-$ or $\gamma \rightarrow e^+e^-$	$0.01 \pm 0.004$
$\pi^-$ hard bremsstrahlung in the WCAL, $\gamma \rightarrow e^+e^-$	$< 0.0001$
$\pi, K \rightarrow e\nu, K_{e4}$ decays	$< 0.001$
$eZ \rightarrow eZ\mu^+\mu^-; \mu^\pm \rightarrow e^\pm\nu\nu$	$< 0.001$
punchthrough $\pi$	$< 0.003$
Total	$0.07 \pm 0.035$




# Results from 2017 run, $5.4 \times 10^{10}$ EOT



# Results from Sept'2017 run, $5.4 \times 10^{10}$ EOT



NA64 exclusion area,  $5.4 \times 10^{10}$  EOT



$X$  is simulated as  $A'$  in invisible mode,  
then decayed with  
 $\Gamma \sim m\epsilon^2$ ,  $\text{Br}(X \rightarrow e^+e^-)=1$

Part of the  ${}^8\text{Be}^*$  region (red vertical line)  
is excluded:  $1.3 \times 10^{-4} < \epsilon_e < 4.2 \times 10^{-4}$

Region of  $A'$  with different masses  
decaing to  $e^+e^-$  is excluded


arXiv:1803.07748 [hep-ph],  
published in Phys. Rev. Letters




# Further running and expected results

Some more data (less than expected) were taken in June 2018 with the visible mode configuration optimized for bigger  $\varepsilon$ . NA64 running after LHC LS2 (in 2021) is preliminarily approved.

Exclusion area for  $A' \rightarrow e^+e^-$  vs EOT



# Conclusion



- A search is performed for sub-GeV dark photons ( $A'$ ) mediated production of dark matter by the NA64 experiment with  $4.3 \cdot 10^{10}$  100 GeV electrons on target
- No evidence for such events is found. This allows to derive an upper limit on the  $A' - \gamma$  mixing strength in the  $A'$  mass range from 1 to 500 MeV and allows to exclude a vector mediator particle solution (universal or  $e$ -coupled) to the  $(g-2)$  anomaly
- More data were taken in 2017 and 2018, are being analysed
- A search is performed for a new X-boson decaying to  $e^+e^-$
- No evidence for such particles are found. This allows to exclude part of the  ${}^8\text{Be}^*$  preferred region and a region on the  $m - \varepsilon$  plane for similar particles with different masses
- Some more data in this configuration were taken in 2018, to be analysed
- The project of running in the muon beams together with COMPASS is being prepared, the purpose is to probe  $(g-2)_\mu$ ,  $\mu - \tau$  conversion etc.



# Backup slides

Old figure

Exclusion area for  $A' \rightarrow e^+e^-$  vs EOT

