

Synchrotron radiation tagging of 100 GeV electrons in NA64 experiment at CERN SPS

Emilio Depero

On behalf of the NA64 collaboration


Institute for Particle Physics, ETH Zürich

5th Beam Telescope 2017 and test Beam result Workshop
January 26, 2017, Barcelona

- 1 Introduction
- 2 Synchrotron radiation
- 3 Results during test beam
- 4 Conclusion

Dark Matter

- Several cosmological observations suggest the existence of Dark Matter, e.g. galaxy rotation curves¹, gravitational lensing and spectrum of CMB radiation
- In addition to the most popular model (WIMP, SUSY etc.) various DM model motivate “light” new physics that could be observed in low energy experiment.
- One possibility is a dark sector of SM singlets coupled to ordinary matter by gravity and possibly other very weak forces. An extra (broken) $U(1)'$ symmetry would implies a new massive boson A' (**Dark Photon**)




Gravitational lensing, from Hubble mission

Dark Matter

- One possibility is a dark sector of SM singlets coupled to ordinary matter by gravity and possibly other very weak forces. An extra (broken) $U(1)'$ symmetry would imply a new massive boson A' (**Dark Photon**)

Dark Photon

- $\Delta L = \epsilon F^{\mu\nu} A'_{\mu\nu}$ kinetic mixing
 γ - A' , ϵ coupling strength
- natural coupling $\epsilon \sim 10^{-3} - 10^{-4}$,
 $M_{A'} \sim \epsilon^{1/2} M_Z$
- Decay Modes:
 - 1 **Visible:** $A' \rightarrow e^- e^+, \mu^- \mu^+$
 - 2 **Invisible:** $A' \rightarrow \chi \bar{\chi}$



Dark Matter

- One possibility is a dark sector of SM singlets coupled to ordinary matter by gravity and possibly other very weak forces. An extra (broken) $U(1)'$ symmetry would imply a new massive boson A' (**Dark Photon**)

Dark Photon


- $\Delta L = \epsilon F^{\mu\nu} A'_{\mu\nu}$ kinetic mixing
 γ - A' , ϵ coupling strength
- natural coupling $\epsilon \sim 10^{-3} - 10^{-4}$,
 $M_{A'} \sim \epsilon^{1/2} M_Z$
- Decay Modes:
 - 1 **Visible:** $A' \rightarrow e^- e^+, \mu^- \mu^+$
 - 2 **Invisible:** $A' \rightarrow \chi \bar{\chi}$ **NA64 focus** \implies
Searching for missing momentum

Motivations

- 1 if $M_{A'} \geq 2M_{DM}$ and $\alpha_{DM} \gg \epsilon^2 \alpha$ invisible decay is the most relevant .
- 2 can explain $(g-2)_\mu$ hint²
- 3 can explain various cosmological observation³

H.S. Lee., Phys. Rev. D 90, 091702 (2014)

N. Arkani-Hamed, D. P. Finkbeiner, T. R. Slatyer, and N. Weiner, Phys. Rev. D 79, 015014 (2009)




Dark photon signal

Dark photon is produced through kinetic mixing of a Bremsstrahlung- γ in the ECAL. The Dark photon would escape then the setup undetected through the invisible decay $A' \rightarrow \chi \bar{\chi}$


Dark Photon signal:

- Missing energy in the ECAL ($E_{ECAL} \leq E_{thresh}$)
- no activity in VETO and HCAL
- reconstructed momentum compatible with a 100 GeV particle
- emitted synchrotron radiation compatible with an electron

Standard model



With Dark photon




Dark photon signal

Dark photon is produced through kinetic mixing of a Bremsstrahlung- γ in the ECAL. The Dark photon would escape then the setup undetected through the invisible decay $A' \rightarrow \chi\bar{\chi}$


Dark Photon signal:


- Missing energy in the ECAL ($E_{ECAL} \leq E_{thresh}$)
- no activity in VETO and HCAL
- reconstructed momentum compatible with a 100 GeV particle
- emitted synchrotron radiation compatible with an electron



NA64 setup

Key features of the experiment:







NA64 setup

Key features of the experiment:

- **High energy particle to trigger the reaction:** 100 GeV e^- from CERN SPS
- **Main impurities of H4 beam:** π^- , low energy e^- ($\sim 1\%$)
 μ^- and K^- ($\leq 0.1\%$)

Accelerator chain of CERN (operating or approved projects)






NA64 setup

Key features of the experiment:

- **High hermeticity:** ECAL PbSc sandwich, $38 \times 38 \times 445 \text{ mm}^3$ ($\sim 40 X_0$) with WLS fiber inserted in spiral and $9\% (E[\text{GeV}])^{-1/2}$ energy resolution






NA64 setup

Key features of the experiment:

- **High hermeticity:** 4 HCAL FeSc sandwich modules, $60 \times 60 \times 150 \text{ cm}^3$ ($\sim 7 \lambda$ for each module) with WLS fiber and $60\% (E[\text{GeV}])^{-1/2}$ energy resolution







NA64 setup

Key features of the experiment:

- Measure momentum:** Tracking system made of 4 MicroMegas modules and 2 GEM detectors together with 2 MPBL magnet [7 T·m] to measure momentum of incoming particles.



Focus of the talk



NA64 setup

Key features of the experiment:

- **Suppress hadronic background:** Synchrotron radiation tagging system (BGO scintillator) to reject μ^- , π^- and K^- decay in flight after interaction with ECAL **Focus of the talk**



- 1 Introduction
- 2 **Synchrotron radiation**
- 3 Results during test beam
- 4 Conclusion

Synchrotron radiation is the radiation emitted by a charged particle accelerated radially to its momentum's direction.

The radiated power emitted per meter can be expressed as:

$$P = \frac{e^2 c}{6\pi_0 (mc^2)^4} \frac{E^4}{R^2} \quad (1)$$

- Power emitted scales on the fourth power of the **energy**. \Rightarrow works better for high e^- momentum
- Power emitted scales to the inverse fourth power of **mass** \Rightarrow very high separation between heavy charged particle and e^- ! (basically no radiation below 1 TeV for particles heavier than e^-)
- Scales as the inverse square power of the radius of the particle track \Rightarrow Total emitted power is proportional to the integrated field seen by the particle.
- emission angle $\theta_{synch} \sim 1/\gamma$ in the laboratory system, basically collinear to particle's momentum at high energies.


In **NA64** heavy charged particle (mainly π^- and μ^-) can leave a fake signal with invisible decay or neutral punchthrough.

⇒ They can be rejected by measuring emitted synchrotron radiation during their passage in the MPBL magnets.

In **NA64** heavy charged particle (mainly π^- and μ^-) can leave a fake signal with invisible decay or neutral punchthrough.

⇒ They can be rejected by measuring emitted synchrotron radiation during their passage in the MPBL magnets.

- Synchrotron photon are detected with two row of 4 BGO crystal placed in the ~ 32 cm space between the deflected and undeflected beam
- First two crystal of each row are used to detect synchrotron radiation (**SRD BGO**). Last two crystal in each row are used to reject hard bremsstrahlung events and backscattering from ECAL (**VETO BGO**).



In **NA64** heavy charged particle (mainly π^- and μ^-) can leave a fake signal with invisible decay or neutral punchthrough.

⇒ They can be rejected by measuring emitted synchrotron radiation during their passage in the MPBL magnets.

BGO detector

- 2 rows of 4 BGO hexagonal crystal, 55mm external diameter and 200mm length
- Wrapped in Teflon tape to improve light collection efficiency
- glued to EMI 9603 PMT
- Very high probability of photoelectric absorption due to high Z
- 8500 γ /MeV light yield
- energy resolution of $\sim 17\%$ (FWHM) at 1.27 MeV
- Decay time of ~ 300 ns

In **NA64** heavy charged particle (mainly π^- and μ^-) can leave a fake signal with invisible decay or neutral punchthrough.


⇒ They can be rejected by measuring emitted synchrotron radiation during their passage in the MPBL magnets.

- Method was previously tried for smaller energies⁴(~ 50 GeV), with smaller efficiency(77%) and suppression factor (10^{-2})
- Good alternative to Cherenkov at high energies for e^- tagging⁵:
 - Energy radiated is higher
 - emission angle $\theta_{cher} \propto 1/\beta$ and at high energy $|\beta_{\pi^-} - \beta_{e^-}|$ is very small
- Requires relatively simple detector (scintillator)

^dW.Busza, M.Chen and D.Luckey, Application of synchrotron radiation for the identification of electrons in high energy beam, Nuclear instrument and method in physics, 105(1972)


^eE. Nappi, RICH detector, CERN EP/99-149, 1999

- The suppression factor for particle heavier than e^- should be in the order of $(m_e/m_{\pi^-})^4 \leq 10^{-8}$ only considering synchrotron radiation. However MC simulations show larger fraction of high energy deposit.



K.A. Olive et al. (PDG), Passage of particles through matter, Chin. Phys., C38, 2014

- Most of the energy lost by heavy particle with energy in the MIP range is due low energy electron ionization (typically few keV), rare high-energy electron ionization interactions (**knock-on** electrons) however can leave enough energy to mimic a synchrotron signal and limit the suppression to $\sim 10^{-3}$ using simple cut on total energy deposited in the SRD detector.



- Most of the energy lost by heavy particle with energy in the MIP range is due low energy electron ionization (typically few keV), rare high-energy electron ionization interactions (**knock-on** electrons) however can leave enough energy to mimic a synchrotron signal and limit the suppression to $\sim 10^{-3}$ using simple cut on total energy deposited in the SRD detector.


Distribution of electrons with kinetic energy T much larger than ionization energy I ($T \gg I$) can be described as⁶:

$$\frac{dN}{dTdx} = \frac{1}{2} K z^2 \frac{Z}{A} \frac{1}{\beta^2} \frac{F(T)}{T^2} \quad (2)$$

- Number N of electron emitted with energy T roughly dependent on the inverse of T
- $T \leq W_{max}$, where W_{max} is the maximum energy transfer for a particle to an electron which scales roughly as the square of the particle energy for heavy particle ($W_{max} \propto \gamma^2$)


K.A. Olive et al. (PDG), Passage of particles through matter, Chin. Phys., C38, 2014

- Most of the energy lost by heavy particle with energy in the MIP range is due low energy electron ionization (typically few keV), rare high-energy electron ionization interactions (**knock-on** electrons) however can leave enough energy to mimic a synchrotron signal and limit the suppression to $\sim 10^{-3}$ using simple cut on total energy deposited in the SRD detector.
- **Idea:** Typically no more than one such interaction is produced per event. It should be possible to exploit detector **transverse segmentation** to distinguish synchrotron radiation (homogeneously distributed in the whole detector) from knock-on electrons (spacially localized since most of the energy is transported by a single electron).




- 1 Introduction
- 2 Synchrotron radiation
- 3 Results during test beam**
- 4 Conclusion


- This Method was tested during the beam time in July 2016 with the BGO detector, the ability of rejecting π^- and μ^- efficiently was confirmed both for the nominal 100 GeV e^- of the experiment using ECAL and HCAL to distinguish incoming particles and directly using a 100 GeV π^- beam ($\sim 9.8 \times 10^4$ events collected).
- A detailed MC simulation of the full **NA64** setup was prepared and the results was compared to the acquired data.




- **Results:** generally very good agreement between the spectra for typical synchrotron energies.
- Some disagreement in the high energy spectrum of electron were recovered by eliminating high energy contamination using the **VETO BGO** and by adding the expected pileup ($\sim 1\%$) for the intensity of the beam ($\sim 3.5 \times 10^5$ particles/spill)





- **Results:** generally very good agreement between the spectra for typical synchrotron energies.
- some disagreement in synchrotron range between data and simulation for π^- . Probably due beam impurities.




- Number of crystal **activated** (more than 1 MeV of energy deposition) was checked in the data to be correlated with particle type as expected.
- few events in the π^- run with 2 activated crystals were eliminated by requiring the total energy deposited in the BGO detector to be larger than 5 MeV. Such events are probably caused by knock-on electrons that further ionize during their travel to the detector.








event accepted from BGO for a cut of $1 < E < 500$ MeV



event rejected from BGO for a cut of 1 MeV



- 1 Introduction
- 2 Synchrotron radiation
- 3 Results during test beam
- 4 Conclusion

- **Synchrotron radiation** can be used to efficiently tag electrons in a beam by placing a **scintillator** in the region between undeflected and deflected beam.
- **transverse segmentation** of the detector can be exploited to detect spacial distribution of the synchrotron radiation and improve further the tagging, bringing the suppression factor down to $\sim 10^{-5}$ and allowing so to suppress μ^- and hadrons down to $\leq 10^{-12}$ per e^- in the **NA64** setup.
- efficiency above 95%


- **Synchrotron radiation** can be used to efficiently tag electrons in a beam by placing a **scintillator** in the region between undeflected and deflected beam.
- **transverse segmentation** of the detector can be exploited to detect spacial distribution of the synchrotron radiation and improve further the tagging, bringing the suppression factor down to $\sim 10^{-5}$ and allowing so to suppress μ^- and hadrons down to $\leq 10^{-12}$ per e^- in the **NA64** setup.
- efficiency above 95%

Upgrade:

- some faster scintillator can help reduce pileup at high intensity (PbSc sandwich with decay time $20\text{ns} \leq$ and larger segmentation has been tested), green extended R9420-100 hamamatsu PMT to improve energy resolution and timing.
- larger segmentation might be helpful but smaller crystal means lower threshold \Rightarrow Energy resolution might become an issue

Result of July test beam:

- collected $\sim 2.75 \times 10^9$ events on target.
- 90% exclusion plot already covers a good portion of unexplored parameter space!⁷
- collected $> 10^{10}$ events on target during October 2016 test run \rightarrow analysis in progress...



Search for Invisible Decays of Sub-GeV Dark Photons in Missing-Energy Events at the CERN SPS, Phys. Rev.

Lett. 118,011802(2017)

Emilio Depero (IPP – ETHZ)

synchrotron tagging


5th Beam Telescope Workshop 2017

15 / 16

Thank You!

Questions?

5 Back-Up Slides



efficiency vs purity for different energy resolution

